

## **PARTICULARS TO APPEAR ON THE OUTER PACKAGE**

### **1. NAME OF THE VETERINARY MEDICINAL PRODUCT**

Ventipulmin Syrup 25 micrograms/ml

### **2. STATEMENT OF ACTIVE AND OTHER SUBSTANCES**

Composition:

1 ml Ventipulmin® Syrup contains 25 micrograms clenbuterol hydrochloride plus 1.8 mg methyl parahydroxybenzoate E218 and 0.2 mg propyl parahydroxybenzoate E214 as preservatives.

### **3. PHARMACEUTICAL FORM**

### **4. PACKAGE SIZE**

355 ml

### **5. TARGET SPECIES**

### **6. INDICATION(S)**

Indications:

Treatment of respiratory disease in horses where airway obstruction due to bronchospasm and/or accumulation of mucus is a contributing factor and improved mucociliary clearance is desirable. To be used alone or as adjuvant therapy. (See package leaflet).

### **7. METHOD AND ROUTE(S) OF ADMINISTRATION**

Administration and Dosage:

For oral use. One complete depression of plunger delivers 4 ml syrup. Add 4 ml of syrup per 125 kg body weight to feed, twice daily. Thus, an adult horse weighing 500 kg (approx. 1000 lbs) should receive 16 ml of syrup i.e. 4 complete depressions of the plunger, twice daily.

### **8. WITHDRAWAL PERIOD**

Withdrawal periods:

Horses intended for human consumption should not be slaughtered until 28 days after treatment.

Do not use in animals producing milk for human consumption.

### **9. SPECIAL WARNING(S), IF NECESSARY**

**Warnings:**

Ventipulmin® antagonises the effects of prostaglandin F2 alpha and oxytocin.

Ventipulmin® is antagonised by beta-adrenergic blocking agents. If used during pregnancy, treatment must be discontinued at the expected time of delivery since uterine contractions may be abolished under its influence. This product contains clenbuterol, a beta-agonist. Take care to avoid skin contact. In case of skin contact wash affected area thoroughly. If irritation occurs/persists seek medical advice. Take care to avoid accidental eye contact. In the case of accidental eye contact, flush thoroughly with clean water and seek medical advice. When using do not eat, drink or smoke. Wash hands thoroughly after using the product. For further contra-indications and warnings, see package leaflet.

## **10. EXPIRY DATE**

Expiry Date:

## **11. SPECIAL STORAGE CONDITIONS**

Do not store above 25°C. Protect from light. Keep bottle in outer carton.  
This pack should be used within 30 days of first opening. Discard unused material.

## **12. SPECIFIC PRECAUTIONS FOR THE DISPOSAL OF UNUSED PRODUCTS OR WASTE MATERIALS, IF ANY**

Dispose of any unused product and empty containers in accordance with guidance from your local waste regulation authority.

## **13. THE WORDS “FOR ANIMAL TREATMENT ONLY” AND CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE IF APPLICABLE**

For animal treatment only.  
To be used in accordance with the directions of a veterinary surgeon.

VM 08327/5029 : POM-V : To be supplied only on veterinary prescription.
---

## **14. THE WORDS “KEEP OUT OF THE SIGHT AND REACH OF CHILDREN”**

Keep out of reach and sight of children.

## **15. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER**

Marketing Authorisation Holder  
Boehringer Ingelheim Animal Health UK Limited, Ellesfield Avenue, Bracknell,  
Berkshire, RG12 8YS, United Kingdom

**16. MARKETING AUTHORISATION NUMBER(S)**

Vm 08327/5029

**17. MANUFACTURER'S BATCH NUMBER**

Batch No.:

**18. OTHER INFORMATION**

Veterinary Bronchodilator

## **MINIMUM PARTICULARS TO APPEAR ON SMALL IMMEDIATE PACKAGING UNITS**

### **1. NAME OF THE VETERINARY MEDICINAL PRODUCT**

Ventipulmin Syrup 25 micrograms/ml

### **2. QUANTITY OF THE ACTIVE SUBSTANCE(S)**

Composition: 1 ml Ventipulmin® Syrup contains 25 micrograms clenbuterol hydrochloride plus 1.8 mg methyl parahydroxybenzoate E218 and 0.2 mg propylhydroxybenzoate E214 as preservatives.

### **3. CONTENTS BY WEIGHT, BY VOLUME OR BY NUMBER OF DOSES**

355 ml

### **4. ROUTE(S) OF ADMINISTRATION**

Administration and Dosage: For oral use. One complete depression of plunger delivers 4 ml syrup. Add 4 ml of syrup per 125 kg body weight to feed, twice daily. Thus, an adult horse weighing 500 kg (approx. 1000 lbs) should receive 16 ml of syrup i.e. 4 complete depressions of the plunger, twice daily.

### **5. WITHDRAWAL PERIOD**

Withdrawal periods: Horses intended for human consumption should not be slaughtered until 28 days after treatment.

Do not use in animals producing milk for human consumption.

### **6. BATCH NUMBER**

Batch No.:

### **7. EXPIRY DATE**

Expiry Date:

### **8. THE WORDS “FOR ANIMAL TREATMENT ONLY”**

For animal treatment only.

### **9. OTHER INFORMATION**

Indications: Treatment of respiratory disease in horses where airway obstruction due to bronchospasm and/or accumulation of mucus is a contributing factor and improved mucociliary clearance is desirable. To be used alone or as adjuvant therapy.

(See package leaflet).

Warnings: Ventipulmin® antagonises the effects of prostaglandin F2 alpha and oxytocin.

Ventipulmin® is antagonised by betaadrenergic blocking agents. If used during pregnancy, treatment must be discontinued at the expected time of delivery since uterine contractions may be abolished under its influence.

This product contains clenbuterol, a beta-agonist. Take care to avoid skin contact. In case of skin contact wash affected area thoroughly. If irritation occurs/persists seek medical advice. Take care to avoid accidental eye contact. In case of accidental eye contact, flush thoroughly with clean water and seek medical advice. When using do not eat, drink or smoke. Wash hands thoroughly after using the product. For further contra-indications and warnings, see package leaflet.

Do not store above 25°C. Protect from light. This pack should be used within 30 days of first opening. Discard unused material. Dispose of any unused product and empty containers in accordance with guidance from your local waste regulation authority.

Keep out of reach and sight of children.

Keep bottle in outer carton.

To be used in accordance with the directions of a veterinary surgeon. To be supplied only on veterinary prescription.

VM 08327/5029 : POM-V

Marketing Authorisation Holder  
Boehringer Ingelheim Animal Health UK Limited, Ellesfield Avenue, Bracknell,  
Berkshire, RG12 8YS, United Kingdom

Veterinary bronchodilator

**[Include information under these headings as it appears in the SPC]**

**PACKAGE LEAFLET FOR: Ventipulmin Syrup 25 micrograms/ml Clenbuterol hydrochloride**

**1. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER AND OF THE MANUFACTURING AUTHORISATION HOLDER RESPONSIBLE FOR BATCH RELEASE, IF DIFFERENT**

**Marketing authorization holder**

Boehringer Ingelheim Animal Health UK Limited, Ellesfield Avenue, Bracknell, Berkshire, RG12 8YS, United Kingdom

**Manufacturer for batch release:**

Boehringer Ingelheim Vetmedica GmbH,  
D 55216, Ingelheim am/Rhein,  
Germany.

**2. NAME OF THE VETERINARY MEDICINAL PRODUCT**

Ventipulmin syrup 25 micrograms/ml

**3. STATEMENT OF THE ACTIVE SUBSTANCE (S) AND OTHER INGREDIENTS**

**Presentation:**

A clear colourless syrup. Each ml contains 25 micrograms clenbuterol hydrochloride plus 1.8 mg methylparahydroxybenzoate E218 and 0.2 mg propyl parahydroxybenzoate E214 as preservatives.

**4. INDICATION(S)**

**Uses:**

Treatment of respiratory disease in horses where airway obstruction due to bronchospasm and/or accumulation of mucus is a contributing factor and improved mucociliary clearance is desirable. To be used alone or as adjuvant therapy. In cases accompanied by bacterial infection the administration of antimicrobial agents is recommended.

**In particular:**

1. Acute, sub-acute and chronic infections where the presence of mucus and/or micro-organisms may stimulate bronchospasm or cause airway obstruction and thus increase airway resistance. For example, bronchitis, bronchiolitis and bronchopneumonia alone or associated with equine influenza and other viral respiratory diseases.
2. Acute, sub-acute and chronic respiratory allergies.
3. Chronic obstructive pulmonary disease (COPD) in horses.

## **5. CONTRAINDICATIONS**

Contra-indications, warnings etc:

Do not use in cases of known hypersensitivity to clenbuterol. Do not use in horses with known cardiac disease.

Ventipulmin® antagonises the effects of prostaglandin F2-alpha and oxytocin and is antagonised by  $\beta$ -adrenergic blocking agents.

In case of accidental overdose, a  $\beta$ - blocker such as propranolol may be used as an antidote.

If used during pregnancy, treatment must be discontinued at the expected time of delivery since uterine contractions may be abolished under its influence.

## **6. ADVERSE REACTIONS**

Adverse effects

Clenbuterol may cause side-effects such as sweating (mainly neck region), muscle tremor, tachycardia, slight hypotension or restlessness. These are typical for  $\beta$ -agonists and occur rarely. If you notice any serious effects or other effects not mentioned in this leaflet, please inform your veterinary surgeon.

## **7. TARGET SPECIES**

## **8. DOSAGE FOR EACH SPECIES, ROUTE(S) AND METHOD OF ADMINISTRATION**

Dosage and Administration:

For oral use.

Dosage: Twice daily administration of 0.8 micrograms of clenbuterol hydrochloride per kg bodyweight. This corresponds to a twice daily administration of 4ml of the syrup per 125kg bodyweight. (One depression of the pump delivers 4 ml syrup). The syrup should be added to the feed. Treatment should be continued for as long as necessary. To be supplied only on veterinary prescription.

## **9. ADVICE ON CORRECT ADMINISTRATION**

## **10. WITHDRAWAL PERIOD(S)**

Withdrawal Periods

Horses intended for human consumption should not be slaughtered until 28 days after treatment.

Do not use in animals producing milk for human consumption.

## **11. SPECIAL STORAGE PRECAUTIONS**

Pharmaceutical Precautions:

Do not store above 25°C. Protect from light.

This pack should be used within 30 days of opening. Discard unused material. Do not use after the expiry date stated on the label and carton.

## 12. SPECIAL WARNING(S)

User precautions

This product contains clenbuterol, a  $\beta$ -agonist. Take care to avoid skin contact. In case of skin contact wash affected area thoroughly. If irritation occurs/persists seek medical advice. Take care to avoid accidental eye contact. In case of accidental eye contact, flush thoroughly with clean water and seek medical advice. When using do not eat, drink or smoke. Wash hands thoroughly after using the product.

## 13. SPECIAL PRECAUTIONS FOR THE DISPOSAL OF UNUSED PRODUCT OR WASTE MATERIALS, IF ANY

Disposal Advice:

Dispose of any unused product and empty containers in accordance with guidance from your local waste regulation authority.

## 14. PID LINK (Do not print heading)

*[The following statement must be included where reference to the European Union Product Database is included on the product information. This statement is relevant to both UK(GB) and UK(NI) products:]*

Find more product information by searching for the 'Product Information Database' on [www.gov.uk](http://www.gov.uk).

## 15. OTHER INFORMATION

Veterinary Bronchodilator

Package Quantities:

Screw top polyethylene bottle containing 355 ml syrup with 4 ml pump dispenser.

Marketing Authorisation Numbers

VM 08327/5029 : POM-V : To be supplied only on veterinary prescription.
--

FOR ANIMAL TREATMENT ONLY.

KEEP OUT OF REACH AND SIGHT OF CHILDREN.

*Gavin Hall*  
Approved: 29 September 2025